



(19)

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 369 472 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
10.12.2003 Bulletin 2003/50

(51) Int Cl. 7: C11D 3/39

(21) Application number: 03076383.3

(22) Date of filing: 08.05.2003

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PT RO SE SI SK TR
Designated Extension States:
AL LT LV MK

(30) Priority: 06.06.2002 GB 0212984

(71) Applicants:
• UNILEVER N.V.
3013 AL Rotterdam (NL)
Designated Contracting States:
AT BE BG CH LI CZ DE DK EE ES FI FR GR HU
IT LU MC NL PT SE SK TR
• UNILEVER PLC
London EC4P 4BQ (GB)
Designated Contracting States:
GB IE

(72) Inventors:
• Appel, Adrianus Cornelis Maria
3133 AT Vlaardingen (NL)
• Hage, Ronald
3133 AT Vlaardingen (NL)
• Hermant, Roelant Mathijs
3133 AT Vlaardingen (NL)
• Lienke, Joachim
3133 AT Vlaardingen (NL)
• Veerman, Simon Marinus
3133 AT Vlaardingen (NL)

(74) Representative: Elliott, Peter William et al
Unilever plc
Patent Division
Colworth House
Sharnbrook
Bedford MK44 1LQ (GB)

(54) **Preserved enhancement of bleaching catalyst**

(57) The present invention provides a catalytic bleaching system that is low in surfactant alkyl hydroperoxide and comprises an antioxidant.

EP 1 369 472 A1

Description

FIELD OF INVENTION

- 5 [0001] This invention relates to the enhancement of bleaching compositions that are substantially devoid of peroxy species.

BACKGROUND OF INVENTION

- 10 [0002] The use of bleaching catalysts for stain removal has been developed over recent years. The recent discovery that some catalysts are capable of bleaching effectively in the absence of an added peroxy source has recently become the focus of some interest, for example: WO8965905; WO0012667; WO0012808; WO0029537, and, WO0060045.
- [0003] Ways of enhancing the activity or improving the stain bleaching profile of these catalysts are desired.

SUMMARY OF INVENTION

- [0004] The bleaching of a stain by a peroxy species is aided by the presence of an active transition metal catalyst. A peroxy species commonly found in laundry bleaching compositions is hydrogen peroxide (H_2O_2) or a precursor thereof, e.g., sodium percarbonate or sodium perborate. In many instances an activator/precursor, e.g., TAED (tetraacetyliethylene diamine), is present which serves together with hydrogen peroxide to form a peracid $[RC(O)OOH]$ to facilitate bleaching.

- 20 [0005] Recently we have found that oily stains are bleached in the presence of selected transition metal catalysts in the absence of an added peroxy source. The bleaching of an oily stain in the absence of an added peroxy source has been attributed to oxygen derived from the air. Whilst it is true that bleaching is effected by oxygen sourced from the air the route in which oxygen plays a part is becoming understood.

- [0006] We have concluded from our research that bleaching of a chromophore in an oily stain is effected by products formed by adventitious oxidation of components in the oily stain. These products, alkyl hydroperoxides, are generated naturally by autooxidation of the oily stain and the alkyl hydroperoxides together with a transition metal catalyst serve to bleach chromophores in the oily stain. Alkyl hydroperoxides (ROOH) are generally less reactive than other peroxy species, for example, peracids $RC(O)OOH$, hydrogen peroxide (H_2O_2), percarbonates and perborates.

- 30 [0007] A surfactant will migrate to a hydrophilic/hydrophobic interface in an aqueous washing medium. In this regard, because the catalysts serve to bleach with alkyl hydroperoxides found in a stain, a consequence of keeping the surfactant as low as possible in alkyl hydroperoxide content is that bleaching is focused on the oily stain where the alkyl hydroperoxides occur and bleaching is minimised elsewhere.

- 35 [0008] The low level of surfactant hydroperoxide also serves to preserve the integrity of the transition metal catalyst. In addition, by keeping the alkyl hydroperoxide level low, peroxide-catalyst interaction is reduced when the composition is in the form of a liquid, be it in a wash or as a commercial liquid formulation.

- [0009] A problem with the detergent composition is that of maintaining the low level of surfactant HPO or reducing further autooxidation of the surfactant present during storage. The presence of an antioxidant in the detergent composition reduces further autooxidation of the surfactant present during storage.

- 40 [0010] The present invention provides a bleaching composition comprising at least 0.0001 % w/wt of an antioxidant, 1 to 60 % w/wt of a surfactant and an organic substance which forms a complex with a transition metal for bleaching an oily stain the bleaching composition upon addition to an aqueous medium providing an aqueous bleaching medium substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-generating bleach system, the total surfactant contribution with a HLB greater than 10 having a hydroperoxide content (HPO) of less than 100 mMol/Kg, preferably less than 50 mMol/Kg, most preferably less than 15 mMol/Kg.

- 45 [0011] The peroxide level of surfactant having a surfactant-peroxide level is expressed in mMol of hydroperoxide ($-OOH$) present per Kg.

- 50 [0012] The term "substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-generating bleach system" should be construed within spirit of the invention. It is preferred that the composition has as low a content of peroxy species present as possible. It is preferred that the bleaching formulation contains less than 1 % w/wt total concentration of peracid or hydrogen peroxide or source thereof, preferably the bleaching formulation contains less than 0.3 % w/wt total concentration of peracid or hydrogen peroxide or source thereof, most preferably the bleaching composition is devoid of peracid or hydrogen peroxide or source thereof.

- 55 [0013] The surfactant has an HLB (hydrophilic/lipophilic balance) greater than 5, more preferably greater than 10, and most preferably greater than 15. For a discussion of HLB the reader is directed to and article by Griffin, W. C., in J. Soc. Cosmetic Chemists Vol. 1 page 311, 1945 and Davies, J. T. and Rideal, E. K. in Interfacial Phenomena, Acad. Press, NY, 1961, pages 371 to 382. The HLB value requirement reflects the importance of the rate of solubility and

dispersibility of the surfactant present from the bleaching composition to the aqueous wash medium in conjunction with surface activity towards the substrate being washed. The threshold value of HLB as required excludes compounds that do not have the required surfactant properties, for example linoleic or oleic acid have an HLB of 0.8.

[0014] It is preferred that the bleaching composition comprises at least 10 % wt/wt, preferably 15 %, most preferably 20 %, of the total surfactant contribution having an HLB greater than 10.

[0015] The present invention extends to a method of bleaching a substrate comprising applying to the substrate, in an aqueous medium, the bleaching composition according to the present invention.

[0016] The present invention extends to a commercial package comprising the bleaching composition according to the present invention together with instructions for its use.

[0017] The bleaching composition may be contacted to the textile fabric in any suitable manner. For example, it may be applied in dry form, such as in powder form, or in a liquor that is then dried, for example as an aqueous spray-on fabric treatment fluid or a wash liquor for laundry cleaning, or a non-aqueous dry cleaning fluid or spray-on aerosol fluid.

[0018] Any suitable textile that is susceptible to bleaching or one that one might wish to subject to bleaching may be used. Preferably the textile is a laundry fabric or garment.

[0019] In a preferred embodiment, the method according to the present invention is carried out on a laundry fabric using an aqueous treatment liquor. In particular, the treatment may be effected in a wash cycle for cleaning laundry. More preferably, the treatment is carried out in an aqueous detergent bleach wash liquid.

[0020] In a preferred embodiment, the treated textile is dried, by allowing it to dry under ambient temperature or at elevated temperatures. The elevated temperatures are commonly provided by a heated agitated environment, as for example found in a tumble dryer, which has been found to accelerate and enhance the air bleaching effect. The effect of ironing the treated textile also serves to accelerate bleaching.

[0021] The bleaching method may be carried out by simply leaving the substrate in contact with the bleaching composition for a sufficient period of time. Preferably, however, the bleaching composition is in an aqueous medium, and the aqueous medium on or containing the substrate is agitated.

[0022] The bleaching composition may be contacted with the textile fabric in any conventional manner. For example it may be applied in dry form, such as in powder form, or in a liquor that is then dried, for example in an aqueous spray-on fabric treatment fluid or a wash liquor for laundry cleaning, or a non-aqueous dry cleaning fluid or spray-on aerosol fluid.

[0023] In a particularly preferred embodiment the method according to the present invention is carried out on a laundry fabric using aqueous treatment liquor. In particular the treatment may be effected in, or as an adjunct to, an essentially conventional wash cycle for cleaning laundry. More preferably, the treatment is carried out in an aqueous detergent wash liquor. The bleaching composition can be delivered into the wash liquor from a powder, granule, pellet, tablet, block, bar or other such solid form. The solid form can comprise a carrier, which can be particulate, sheet-like or comprise a three-dimensional object. The carrier can be dispersible or soluble in the wash liquor or may remain substantially intact. In other embodiments, the bleaching composition can be delivered into the wash liquor from a paste, gel or liquid concentrate.

[0024] A unit dose as used herein is a particular amount of the bleaching composition used for a type of wash. The unit dose may be in the form of a defined volume of powder, granules or tablet.


[0025] It is preferred that a unit dose provides at least 0.05 g/l, preferably 0.1 g/l, most preferably 0.2 g/l, concentration of the surfactant compound in a wash. Ideally, a unit dose in an aqueous medium provides aqueous medium having an HPO not greater than 10 μ M.

DETAILED DESCRIPTION OF THE INVENTION

Antioxidant


[0026] The compositions of the present invention comprise an effective amount of the anti-oxidant, at least 0.0001 % wt/wt, preferably from about 0.001 % more preferably from about 0.1%, most preferably from about 0.2% to about 10%, preferably to about 5%, more preferably to about 1% by weight of an anti-oxidant. Anti-oxidants are substances as described in Kirk-Othmers (Vol 3, pg 424) and in Uhlmanns Encyclopedia (Vol 3, pg 91).

[0027] One class of anti-oxidants suitable for use in the present invention is alkylated phenols having the general formula:



wherein R is C1-C22 linear or branched alkyl, preferably methyl or branched C3-C6 alkyl; C3-C6 alkoxy, preferably methoxy; R1 is a C3-C6 branched alkyl, preferably tert-butyl; x is 1 or 2. Hindered phenolic compounds are preferred as antioxidant.

[0028] Another class of anti-oxidants suitable for use in the present invention is a benzofuran or benzopyran derivative having the formula:



wherein R1 and R2 are each independently alkyl or R1 and R2 can be taken together to form a C5-C6 cyclic hydrocarbyl moiety; B is absent or CH2; R4 is C1-C6 alkyl; R5 is hydrogen or -C(O)R3 wherein R3 is hydrogen or C1-C19 alkyl; R6 is C1-C6 alkyl; R7 is hydrogen or C1-C6 alkyl; X is -CH2OH, or -CH2A wherein A is a nitrogen comprising unit, phenyl, or substituted phenyl. Preferred nitrogen comprising A units include amino, pyrrolidino, piperidino, morpholino, piperazino, and mixtures thereof.

[0029] Other suitable antioxidants are found as follows. A derivative of α -tocopherol, 6-hydroxy-2,5,7,8-tetra-methylchroman-2-carboxylic acid (TroloxTM).

Anti-oxidants/radical scavengers such as ascorbic acid (vitamin C) and its salts, tocopherol (vitamin E), tocopherol sorbate, other esters of tocopherol, butylated hydroxy benzoic acids and their salts, gallic acid and its alkyl esters, especially propyl gallate, uric acid and its salts and alkyl esters, sorbic acid and its salts, the ascorbyl esters of fatty acids, amines (e.g., N,N-diethylhydroxylamine, amino-guanidine), sulphydryl compounds (e.g., glutathione), and dihydroxy fumaric acid and its salts may be used.

[0030] Non-limiting examples of anti-oxidants suitable for use in the present invention include phenols *inter alia* 2,6-di-tert-butylphenol, 2,6-di-tert-butyl-4-methylphenol, mixtures of 2 and 3- tert-butyl-4-methoxyphenol, and other ingredients including include propyl gallate, tert-butylhydroquinone, benzoic acid derivatives such as methoxy benzoic acid, methylbenzoic acid, dichloro benzoic acid, dimethyl benzoic acid, 5-hydroxy-2,2,4,6,7-pentamethyl-2,3-dihydro-1-benzofuran-3-one, 5-hydroxy-3-methylene-2,2,4,6,7-pentamethyl-2,3-dihydro-benzofuran, 5-benzyloxy-3-hydroxymethyl-2,2,4,6,7-pentamethyl-2,3-dihydro-1-benzofuran, 3-hydroxymethyl-5-methoxy-2,2,4,6,7-pentamethyl-2,3-dihydro-1-benzofuran, vitamin C(ascorbic acid), and Ethoxyquine (1,2-dihydro-6-ethoxy-2,2,4-trimethylchinolin) marketed under the name RaluquinTM by the company RaschigTM.

[0031] Preferred radical scavengers for use herein include di-tert-butyl hydroxy toluene (BHT), α -tocopherol, hydroquinone, 2,2,4-trimethyl-1,2-dihydroquinoline, di-tert-butyl hydroquinone, mono-tert-butyl hydroquinone, tert-butyl-hydroxy anisole, benzoic acid and derivatives thereof, like alkoxylated benzoic acids, as for example, trimethoxy benzoic acid (TMBA), toluic acid, catechol, t-butyl catechol, benzylamine, 1,1,3-tris(2-methyl-4-hydroxy-5-t-butylphenyl) butane, N-propyl-gallate or mixtures thereof and highly preferred is di-tert-butyl hydroxy toluene.

[0032] It is preferred that at least two antioxidants are present in the composition in a molar ratio of at least 5%, preferably at least 10%, most preferably 25%. The mixture of the two antioxidants providing a greater degree of antioxidant activity than the molar sum of the two antioxidants.

The Surfactant

[0033] Many surfactants are susceptible to autoxidation to form hydroperoxides. Surfactants containing an allylic hydrogen, a hydrogen alpha to an ether or an benzylic hydrogen are particularly susceptible to autoxidation.

[0034] One skilled in the art will appreciate that benzene is considered unsaturated but does not contain allylic hydrogens per se. The homolytic bond dissociation energy (BDE) for benzene (C₆H₅-H) is 110.9 kcal/mol (298 K) makes benzene resistant to autoxidation. An unsaturated compound having a hydrogen atom covalently bound to an alpha-carbon that is alpha to a Sp²-Sp² hybridized bond, e.g., as shown as underlined in the following formula CH₂=CH-CH₂-CH₃, is known as an allylic hydrogen and is susceptible to autoxidation.

Below is a table of bond strengths (298 K) obtained from:

[0035] The handbook of Chemistry and Physics 73rd edition, CRC Press. Generally the lower the C-H bond dissociation energy of a compound the greater the propensity the C-H bond to undergo autoxidation.

Compound	BDE ΔH(kcal/mol)
(CH ₃) ₃ CH	93.3 ± 0.5
H-CH ₂ OCH ₃	93 ± 1
C ₆ H ₅ -H	110.9 ± 2.0
H-CMe ₂ OH	91 ± 1
CH ₃ CH ₃	100.3 ± 1
CH ₂ =CH-CH ₂ -CH ₃	83.1 ± 2.2
CH ₂ =CH-CH ₃	86.3 ± 1.5
C ₆ H ₅ -CH ₃	88.0 ± 1
CH ₃ CH=CHCH=CH ₂	83 ± 3

[0036] It is particularly important to consider the HPO content of surfactant that comprises a hydrogen atom moiety of the following form: -CH=CH-CH₂-, Ar-CH₂-, and/or, -O-CH₂- where the hydrogen atom has a homolytic bond dissociation energy of less than 90 kcal/mol.

[0037] The following is intended as general examples of surfactants that are prone to autoxidation. When used in the present invention the hydroperoxide (HPO) level of these types of surfactants should be examined. In many instances the preparation, storage and handling thereof should be considered to keep autoxidation to a minimum.

[0038] A fatty acid soap used preferably contains from about 16 to about 22 carbon atoms, preferably in a straight chain configuration. Preferably the number of carbon atoms in the fatty acid soap is from about 16 to about 18.

[0039] This soap, in common with other anionic detergents and other anionic materials in the detergent compositions of this invention, has a cation, which renders the soap water-soluble and/or dispersible. Suitable cations include sodium, potassium, ammonium, monethanolammonium, diethanolammonium, triethanolammonium, tetramethylammonium, etc. cations. Sodium ions are preferred although in liquid formulations potassium, monoethanolammonium, diethanolammonium, and triethanolammonium cations are useful.

[0040] The soaps are frequently made from natural oils that often contain one or more unsaturated groups and consist of mixtures of components. It is clear that hydrolysis of these natural components yield mixtures of soaps. Examples of natural oils are sunflower oil, olive oil, cottonseed oil, linseed oil, safflower oil, sesame oil, palm oil, corn oil, peanut oil, soybean oil, castor oil, coconut oil, canola oil, cod liver oil and the like, that give mixtures of soaps. However, also hydrolysis products of purified oils, as listed above, may be employed. Other examples of soaps include erucic acid.

[0041] As one skilled in the art will appreciate a cationic may be manufactured, for example, by adding an alkyl halide to an amine thus forming a cationic.

[0042] In principle the cationic surfactants exhibit the same requirements as listed above for the soap materials, except they need to be quarternised. Without limiting the scope of the invention, suitable cationics may be formed by preparing the quaternary salts from alcohols that were obtained from the corresponding fatty acid. Examples of cationic surfactants based on natural oils include oleylbis(2-hydroxyethyl)trimethylammonium chloride and dilalloy fatty alkylidimethyl ammonium chloride.

[0043] In general, the nonionic and anionic surfactants of the surfactant system may be chosen from the surfactants described "Surface Active Agents" Vol. 1, by Schwartz & Perry, Interscience 1949, Vol. 2 by Schwartz, Perry & Berch,

Interscience 1958, in the current edition of "McCutcheon's Emulsifiers and Detergents" published by Manufacturing Confectioners Company or in "Tenside-Taschenbuch", H. Stache, 2nd Edn., Carl Hauser Verlag, 1981.

[0044] Suitable nonionic detergent compounds which may be used include, in particular, the reaction products of compounds having a hydrophobic group and a reactive hydrogen atom, for example, aliphatic alcohols, acids, amides or alkyl phenols with ethylene oxides, especially ethylene oxide either alone or with propylene oxide. Specific nonionic detergent compounds are C_8 - C_{22} alkyl phenol-ethylene oxide condensates, generally 5 to 25 EO, i.e. 5 to 25 units of ethylene oxide per molecule, and the condensation products of aliphatic C_8 - C_{18} primary or secondary linear or branched alcohols with ethylene oxide, generally 5 to 40 EO.

[0045] Suitable anionic detergent compounds which may be used are usually water-soluble alkali metal salts of organic sulphates and sulphonates having alkyl radicals containing from about 8 to about 22 carbon atoms, the term alkyl being used to include the alkyl portion of higher acyl radicals. Examples of suitable synthetic anionic detergent compounds are sodium and potassium alkyl sulphates, especially those obtained by sulphating higher C_8 - C_{18} alcohols, produced for example from tallow or coconut oil, sodium and potassium alkyl C_8 - C_{20} benzene sulphonates, particularly sodium linear secondary alkyl C_{12} - C_{15} benzene sulphonates, and sodium alkyl glyceryl ether sulphates, especially those ethers of the higher alcohols derived from tallow or coconut oil and synthetic alcohols derived from petroleum. The preferred anionic detergent compounds are sodium C_{11} - C_{18} alkyl benzene sulphonates and sodium C_{12} - C_{18} alkyl sulphates. Also applicable are surfactants such as those described in EPA-328 177 (Unilever), which show resistance to salting-out, the alkyl polyglycoside surfactants described in EPA-070 074, and alkyl monoglycosides.

[0046] Preferred surfactant systems are mixtures of anionic with nonionic detergent active materials, in particular the groups and examples of anionic and nonionic surfactants pointed out in EP-A-346 995 (Unilever). Especially preferred is surfactant system that is a mixture of an alkali metal salt of a C_{16} - C_{18} primary alcohol sulphate together with a C_{12} - C_{15} primary alcohol 3-7 EO ethoxylate. The nonionic detergent is preferably present in amounts greater than 10%.

[0047] The low level of surfactant hydroperoxide also serves to reduce bleaching action towards unwanted substrates in a wash. By reducing the level of hydroperoxides in the bleaching composition, by virtue of low surfactant hydroperoxides, the interaction between the transition metal catalyst and a hydroperoxide is more likely to take place in the stain than elsewhere. In this manner, integrity of an item being washed with the bleaching composition of the present invention is maintained for longer in comparison to when the level of surfactant hydroperoxides is above the criteria given.

The Bleach Catalyst

[0048] The bleach catalyst per se may be selected from a wide range of transition metal complexes of organic molecules (ligands). In typical washing compositions the level of the organic substance is such that the in-use level is from 0.05 μ M to 50 mM, with preferred in-use levels for domestic laundry operations falling in the range 1 to 100 μ M. Higher levels may be desired and applied in industrial textile bleaching processes.

[0049] Suitable organic molecules (ligands) for forming complexes and complexes thereof are found, for example in:

GB 9906474.3; GB 9907714.1; GB 98309168.7, GB 98309169.5;

GB 9027415.0 and GB 9907713.3; DE 19755493; EP 999050;


WO-A-9534628; EP-A-458379; EP 0909809; United States Patent 4,728,455; WO-A-98/39098; WO-A-98/39406, WO 9748787,

WO 0029537; WO 0052124, and WO0060045 the complexes and organic molecule (ligand) precursors of which are herein incorporated by reference. An example of a preferred catalyst is a transition metal complex of MeN4Py ligand (N,N-bis(pyridin-2-yl)-1,1-bis(pyridin-2-yl)-1-aminoethane).

[0050] The ligand forms a complex with one or more transition metals. In the latter case for example as a dinuclear complex. Suitable transition metals include for example: manganese in oxidation states II-IV, iron II-V, copper I-III, cobalt I-III, titanium II-IV, tungsten IV-VI, vanadium II-V and molybdenum II-VI.

[0051] A bleaching composition comprising:

a) a monomer ligand or transition metal catalyst thereof of a ligand having the formula (I):



wherein each R is independently selected from: hydrogen, F, Cl, Br, hydroxyl, C1-C4-alkylo-, -NH-CO-H, -NH-CO-C1-C4-alkyl, -NH2, -NH-C1-C4-alkyl, and C1-C4-alkyl; R1 and R2 are independently selected from: C1-C4-alkyl, C6-C10-aryl, and,

a group containing a heteroatom capable of coordinating to a transition metal, wherein at least one of R1 and R2 is the group containing the heteroatom;

R3 and R4 are independently selected from hydrogen, C1-C8 alkyl, C1-C8-alkyl-O-C1-C8-alkyl, C1-C8-alkyl-O-C6-C10-aryl, C6-C10-aryl, C1-C8-hydroxyalkyl, and $-(CH_2)_nC(O)OR_5$

wherein R5 is independently selected from: hydrogen, C1-C4-alkyl, n is from 0 to 4, and mixtures thereof, and, X is selected from C=O, $-C(R_6)_2-$, wherein Y is from 0 to 3 each R6 is independently selected from hydrogen, hydroxyl, C1-C4-alkoxy and C1-C4-alkyl.

[0052] The transition metal complex preferably is of the general formula (A):



in which:

M represents a metal selected from Mn(II)-(III)-(IV)-(V), Cu(I)-(II)-(III), Fe (II)-(III)-(IV)-(V), Co(I)-(II)-(III), Ti(II)-(III)-(IV), V(II)-(III)-(IV)-(V), Mo(II)-(III)-(IV)-(V)-(VI) and W(IV)-(V)-(VI), preferably from Fe(II)-(III)-(IV)-(V);

L represents the ligand, preferably N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane, or its protonated or deprotonated analogue;

X represents a coordinating species selected from any mono, bi or tri charged anions and any neutral molecules able to coordinate the metal in a mono, bi or tridentate manner;

Y represents any non-coordinated counter ion;

a represents an integer from 1 to 10;

k represents an integer from 1 to 10;

n represents zero or an integer from 1 to 10;

m represents zero or an integer from 1 to 20.

Experimental

Compounds

[0053] The ligand N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane (MeN4py) was prepared as described in EP 0 909 809 A2. $[Fe(MeN4py)Cl]Cl$ was prepared as described in WO 01/16271.

[0054] The transition metal complex dimethyl 2,4-di-(2-pyridyl) -3-methyl-7-(pyridin-2-ylmethyl)-3,7-diaza-bicyclo [3.3.1]nonan-9-one-1,5-dicarboxylate $[N_2Py_3FeCl]Cl$ was prepared as described in WO 02/48301.

Bleaching Experiment

[0055] In an aqueous solution of 6°FH (made by adding 5.45 ml of a stock solution containing 23.5 g/l $CaCl_2 \cdot 2H_2O$ and 8.1 g/l $MgCl_2 \cdot 6H_2O$ to one liter water) containing 2g/L of OMO MA™ Brazil, curry-oil stained cloths were added and kept in contact with the solution whilst agitating for 30 minutes at 30°C.

[0056] The cloths were washed without any additive (blank in table) and 10 μM of $[N_2Py_3FeCl]Cl$ complex without

and with addition of 1 mM of NaLAS-hydroperoxide (NaLAS-HPO; containing 130 mmol HPO per kg NaLAS - ex Albright and Willson).

[0057] After the wash, the cloths were rinsed with water and subsequently dried at 37°C for 2 h and the change in colour was measured with a Linotype-Hell scanner (ex Linotype). The change in colour (including bleaching) is expressed as the ΔE value versus white; a lower ΔE value means a cleaner cloth. The measured colour difference (ΔE) between the washed cloth and the unwashed cloth is defined as follows:

$$\Delta E = [(\Delta L)^2 + (\Delta a)^2 + (\Delta b)^2]^{1/2}$$

wherein ΔL is a measure for the difference in darkness between the washed and unwashed test cloth; Δa and Δb are measures for the difference in redness and yellowness respectively between both cloths. With regard to this colour measurement technique, reference is made to Commission International de l'Eclairage (CIE); Recommendation on Uniform Colour Spaces, colour difference equations, psychometric colour terms, supplement no 2 to CIE Publication, no 15, Colorimetry, Bureau Central de la CIE, Paris 1978. The results are shown below in the tables.

	OMO MA	OMO MA + LAS-HPO
Blank	60	61
[Fe-N2py3Cl]Cl	42	52

[0058] The results presented in the table show that the iron compound bleaches the stain better in absence of surfactant - hydroperoxide (LAS-HPO) than in the presence of LAS-HPO.

Stability Experiment

[0059] The following demonstrates that an antioxidant may be used to reduce the formation of hydroperoxides. The amount of aldehyde formed is reflective of the degree of surfactant autooxidation. A composition containing hydroperoxides is often a dynamic composition in which radical chain reactions are occurring. Hydroperoxides result in the formation of alkoxyl radicals that can undergo beta-scission to form an aldehyde or a ketone. The hydroperoxide levels of the compositions were monitored by examining decomposition products, aldehydes, formed from the alkali metal salt of sodium oleate.

[0060] A liquid formulation was chosen because it reflects the situation that occurs in a granular detergent but over a shorter time scale than that of the granular system.

Base formulation	
component	%
PAS	10%
Nonionic surfactant, ethoxylated fatty alcohol type	18.4%
Oleic acid	10%
Deflocculating polymer, polymer A11 from EP346,995	1%
Silicon oil to control foam	0.03%
KOH	4.1 %
NaOH	0.9%
Citric acid.H2O	5.5%
Glycerol	5%
Borax	1.9%
Anti-dye transfer polymer	0.3%
Protease	0.3%
Lipolase	0.37%

(continued)

Base formulation	
component	%
Amylase	0.15%
Perfume	0.47%

[0061] Liquid formulation A was prepared with 0.03% of $[\text{Fe}(\text{MeN4py})\text{Cl}]\text{Cl}$ by adding 7.5 mg of the solid material in 25 ml liquid formulation A and optionally the anti-oxidant was added (resulting in 0.1%, 0.05% and 0.025%, unless denoted differently, in the formulation respectively). The mixture was stirred vigorously for 10 min and the liquids were then stored at 37°C.

[0062] The anti-oxidants employed were: BHT (2,6-di-*t*-butyl-4-methylphenol), Trolox (6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid, Rauquin (1,2-dihydro-6-ethoxy-2,2,4-trimethylquinolin, vitamin C, Vitamin E (α -tocopherol), vitamin E-acetate (O-acetyl- α -tocopherol), and a mixture of 10% α , 45% δ - and 45% γ -tocopherol. The latter system was 70% pure; the values given in the table are corrected for this purity.

[0063] The results presented in the table below are those of liquid formulations that have been stored at ambient conditions up till 7 weeks. The measurements were made using a Fisons HRGC mega-2-series set-up using a Chrompack CP-SIL 5 CB column (50 m x 0.32 mm, FD 1.2 μm). 3-Hexanone was used as an internal standard. The head-space analysis was done at 70°C.

Amounts of aldehydes detected by GC analysis of the liquid detergent formulations containing catalyst and anti-oxidants.

[0064]

Catalyst	Antioxidant (% in formul.)	Weeks storage	Octanal	Heptanal	Hexanal
-	-	7	0.16	0.05	0.20
+	-	7	0.75	1.16	4.64
+	Rauquin (0.05)	7	0.26	0.08	0.25
+	Rauquin (0.1)	7	0.20	0.07	0.27
+	Trolox (0.05)	7	0.25	0.17	0.95
+	Trolox (0.1)	7	0.27	0.10	0.40
+	Tocopherol-mix (0.17)	7	0.20	0.09	0.67

Claims

1. A bleaching composition comprising at least 0.0001 % wt/wt of an antioxidant, 1 to 60 % wt/wt of a surfactant and an organic substance which forms a complex with a transition metal for bleaching an oily stain the bleaching composition upon addition to an aqueous medium providing an aqueous bleaching medium substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-generating bleach system, the total surfactant contribution with a HLB greater than 10 having a hydroperoxide content (HPO) less than 100 mMol/Kg, preferably less than 50 mMol/Kg, most preferably less than 15 mMol/Kg.
2. A bleaching composition according to claim 1, wherein the antioxidant is present in the composition in the range 0.001 to 10%, preferably from about 0.1% to 10%, and most preferably from 0.2% to 5%.
3. A bleaching composition according to any preceding claim, wherein the antioxidant is selected from the group consisting of: a phenol and an amine.
4. A bleaching composition according to claim 3, wherein the antioxidant is a hindered phenol.
5. A bleaching composition according to any preceding claim, wherein the antioxidant is selected from the group

EP 1 369 472 A1

consisting of: di-tert-butyl hydroxy toluene, Ethoxyquine, α -tocopherol, and 6-hydroxy-2,5,7,8-tetra-methylchroman-2-carboxylic acid.

6. A bleaching composition according to any preceding claim, wherein the bleaching composition comprises at least 20 % wt/wt of the total surfactant contribution having a HLB greater than 10.
7. A bleaching composition according to any preceding claim, wherein the surfactant is selected from the group of cationics.
8. A bleaching composition according to any preceding claim, wherein the surfactant is selected from the group of anionics.
9. A bleaching composition according to any preceding claim, wherein the surfactant is selected from the group of neutral species.
10. A bleaching composition according to claim 8, wherein the surfactant is a linear alkyl benzene sulphonate.
11. A bleaching composition according to any preceding claim, wherein the surfactant comprises an allylic hydrogen atom.
12. A bleaching composition according to any preceding claim, wherein the surfactant comprises a hydrogen atom moiety selected from the following form: $-\text{CH}=\text{CH}-\text{CHR}-$; $\text{Ar}-\text{CHR}-$; and, $-\text{O}-\text{CHR}-$, wherein the hydrogen atom has a homolytic bond dissociation energy of less than 90 kcal/mol and R is a substituent.
13. A bleaching composition according to any preceding claim, wherein the total surfactant contribution having a HLB greater than 10 has a hydroperoxide content of less than 5 mMol/Kg.
14. A bleaching composition according to any preceding claim, wherein the surfactant is present in the composition in an amount such that a unit dose in an aqueous medium provides aqueous medium having an HPO not greater than 10 μM .

European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 03 07 6383

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (INCL.7)
X	WO 95 27774 A (PROCTER & GAMBLE) 19 October 1995 (1995-10-19) * page 13, line 5 - page 17, line 3; claims; examples *	1-14	C11D3/39
P,X	WO 02 072746 A (UNILEVER) 19 September 2002 (2002-09-19) * claims *	1-14	
D,X	WO 98 39406 A (PROCTER & GAMBLE) 11 September 1998 (1998-09-11) * page 87 - page 88 * * page 93 - page 96 * * claims *	1-14	
D,X	WO 00 52124 A (PROCTER & GAMBLE) 8 September 2000 (2000-09-08) * page 4, line 21 - page 15, line 20 * * tables 1, 11 *	1-10, 13, 14 11, 12	
A	* claims *		
D,X	WO 00 60045 A (PROCTER & GAMBLE) 12 October 2000 (2000-10-12) * page 2, line 11 - page 10, line 3 * * example 2; table 1 *	1-10, 13, 14 11, 12	TECHNICAL FIELDS SEARCHED (INCL.7) C11D
A	* claims *		
X	GB 2 309 976 A (PROCTER & GAMBLE) 13 August 1997 (1997-08-13) * page 31; claims *	1	
X	EP 0 690 122 A (PROCTER & GAMBLE) 3 January 1996 (1996-01-03) * page 12, line 1 - page 13, line 44 * * page 18, line 30 - page 20, line 38 * * claims *	1-14	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 1 September 2003	Examiner Serbetsoglou, A
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-technical disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention S : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons * : member of the same patent family, corresponding document</p>			

EP 03 07 6383 (P4/C03)

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 03 07 6383

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

01-09-2003

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9527774	A	19-10-1995	AT 170553 T	15-09-1998
			CA 2187175 A1	19-10-1995
			DE 69504489 D1	08-10-1998
			DE 69504489 T2	20-05-1999
			EP 0754218 A1	22-01-1997
			JP 3009471 B2	14-02-2000
			JP 9511774 T	25-11-1997
			WO 9527774 A1	19-10-1995
			US 5622646 A	22-04-1997
WO 02072746	A	19-09-2002	WO 02072746 A1	19-09-2002
			US 2003054968 A1	20-03-2003
			CA 2431006 A1	19-09-2002
			WO 02072747 A1	19-09-2002
			US 2003054967 A1	20-03-2003
WO 9839406	A	11-09-1998	AU 732147 B2	12-04-2001
			AU 6226298 A	22-09-1998
			BR 9808840 A	04-07-2000
			CN 1260830 T	19-07-2000
			EP 0977828 A1	09-02-2000
			WO 9839406 A1	11-09-1998
			JP 2001513644 T	04-09-2001
			TR 9902673 T2	21-04-2000
			US 2003119698 A1	26-06-2003
			US 2002004473 A1	10-01-2002
			HU 0001927 A2	28-09-2000
			ZA 9801890 A	01-09-1998
WO 0052124	A	08-09-2000	AU 3612700 A	21-09-2000
			BR 0008715 A	02-01-2002
			CA 2364538 A1	08-09-2000
			CN 1342194 T	27-03-2002
			CZ 20013157 A3	17-07-2002
			EP 1157088 A1	28-11-2001
			JP 2002538268 A	12-11-2002
			WO 0052124 A1	08-09-2000
			US 2002193271 A1	19-12-2002
			US 6251701 B1	26-06-2001
WO 0060045	A	12-10-2000	AU 4061900 A	23-10-2000
			WO 0060045 A1	12-10-2000
GB 2309976	A	13-08-1997	BR 9707402 A	06-04-1999
			CA 2245560 A1	14-08-1997
			CN 1214726 A	21-04-1999

EPO FORM 1/2003

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 03 07 6383

This annex lists the patent family members relating to the parent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

01-09-2003

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 2309976 A		EP 0891410 A1	20-01-1999
		JP 3171855 B2	04-06-2001
		JP 11504067 T	06-04-1999
		WO 9729174 A1	14-08-1997
		US 6093343 A	25-07-2000
EP 0690122 A	03-01-1996	GB 2294268 A	24-04-1996
		CA 2151659 A1	31-12-1995
		EP 0690122 A2	03-01-1996
		US 5695679 A	09-12-1997
		AT 204016 T	15-08-2001
		AU 7216594 A	24-01-1995
		DE 69427912 D1	13-09-2001
		DE 69427912 T2	04-04-2002
		EP 0706559 A1	17-04-1996
		US 5698504 A	16-12-1997

EP 03 07 6383

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 369 472 A8

(12)

CORRECTED EUROPEAN PATENT APPLICATION

Note: Bibliography reflects the latest situation

(15) Correction information:

Corrected version no 1 (W1 A1)
INID code(s) 71

(51) Int. Cl.7: **C11D 3/39**

(48) Corrigendum issued on:

31.03.2004 Bulletin 2004/14

(43) Date of publication:

10.12.2003 Bulletin 2003/50

(21) Application number: 03076383.3

(22) Date of filing: 08.05.2003

(84) Designated Contracting States:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PT RO SE SI SK TR
Designated Extension States:
AL LT LV MK

(72) Inventors:

- Appel, Adrianus Cornelis Maria
3133 AT Vlaardingen (NL)
- Hage, Ronald
3133 AT Vlaardingen (NL)
- Hermant, Roelant Mathijs
3133 AT Vlaardingen (NL)
- Lienke, Joachim
3133 AT Vlaardingen (NL)
- Veerman, Simon Marinus
3133 AT Vlaardingen (NL)

(30) Priority: 06.06.2002 GB 0212984

(71) Applicants:

- UNILEVER N.V.
3013 AL Rotterdam (NL)
Designated Contracting States:
AT BE BG CH LI CZ DE DK EE ES FI FR GR HU
IT LU MC NL PT RO SE SI SK TR
- UNILEVER PLC
London EC4P 4BQ (GB)
Designated Contracting States:
CY GB IE

(74) Representative: Elliott, Peter William et al

Unilever plc
Patent Division
Colworth House
Sharnbrook
Bedford MK44 1LQ (GB)

(54) **Preserved enhancement of bleaching catalyst**

(57) The present invention provides a catalytic bleaching system that is low in surfactant alkyl hydroperoxide and comprises an antioxidant.

EP 1 369 472 A8